

### **Thromboangiitis Obliterans - Case Report**

M. Elfarra<sup>1</sup>, D. Rădulescu<sup>2</sup>, I. Peride<sup>2</sup>, A. Niculae<sup>2</sup>, A. Ciocâlțeu<sup>2</sup>, I.A. Checheriță<sup>2</sup>, I. Lascăr<sup>3</sup>, D.R. Sinescu<sup>4</sup>

<sup>1</sup>Department of Vascular Surgery, "St. John" Emergency Clinical Hospital Bucharest, Romania

<sup>2</sup>Department of Nephrology and Dialysis, Clinical Department No. 3, "Carol Davila" University of Medicine and Pharmacy Bucharest

<sup>3</sup>Department of Microsurgery and Plastic Surgery, Emergency Clinical Hospital, Clinical Department No. 11, "Carol Davila" University of Medicine and Pharmacy Bucharest, Romania

<sup>4</sup>Department of Plastic Surgery and Reconstructive Microsurgery, "Elias" Emergency University Hospital, Clinical Department No. 11, "Carol Davila" University of Medicine and Pharmacy Bucharest, Romania

### **Abstract**

Thromboangiitis obliterans (Buerger's disease) represents an inflammatory disease of limbs' small arteries and veins causing vascular thrombosis, and partial or total obstruction. It affects mostly male gender aged < 40 years old. The peculiarity of our case is underlined by presenting a 62 years, chronic tobacco user and not compliant female patient known with thromboangiitis obliterans for almost 15 years. The arteriographic and clinical features with concomitant and sever affected upper and lower limbs are highly suggestive, emphasizing the possibility of Buerger's disease development even in female patients.

**Key words:** thromboangiitis obliterans, clinical features, diagnosis, treatment

Corresponding author: Associated Professor Ionel Alexandru Checherita, MD, PhD  
Department of Nephrology and Dialysis  
Clinical Department No. 3, "Carol Davila"  
University of Medicine and Pharmacy Bucharest  
37th Dionisie Lupu Street, 1st Sector, 020022, Romania  
E-mail: [fizij@yahoo.com](mailto:fizij@yahoo.com); [al.checherita@gmail.com](mailto:al.checherita@gmail.com)