

Cutaneous and Inguinal Lymph Nodes Metastases Disseminated from an Endometrial Serous Carcinoma - Case Report and Literature Review

Laura Rebegea^{1,2}, Ana-Maria Ilie¹, Anca Neagu², Dorel Firescu^{1,3}, Georgiana Bianca Constantin², Rodica Bîrlă^{4,5}, Xenia Bacinschi³, Cristina Șerban^{1,2}

¹"Department of Oncology, Sf. Ap. Andrei" Emergency Clinical Hospital Galati, Romania

²"Dunarea de Jos" University, Faculty of Medicine and Pharmacy, Galati, Romania

³General Surgery II Clinic, "Sf. Ap. Andrei" Emergency Clinical Hospital, Galati, Romania

⁴Center of Excellence in Esophageal Surgery, "Sf. Maria" Clinical Hospital, Bucharest, Romania

⁵Carol Davila University of Medicine and Pharmacy, Bucharest, Romania

Abstract

Cutaneous metastases from endometrial cancer are rare and unusual. This is the case of a 72 years old female patient, diagnosed in 2018 with G3 endometrial serous carcinoma (ESC). At 18 months from the surgical intervention, the patient developed bilateral inguinal lymph nodes metastases and skin secondary lesions; histological and immunohistochemical tests were performed. Computed tomography scan did not indicate visceral secondary lesions, local or regional relapses. We present the treatment practiced, the case's evolution and we discuss about epidemiology, molecular biology, treatment options and management of advanced and local lesions. The appearance of skin metastases in these cases is associated with poor prognosis and treatment options are limited to palliative chemotherapy and radiotherapy.

Key words: serous carcinoma, endometrium, cutaneous metastases